

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

spores are differentiated and set free. The first partial differentiation and movement of the spore-origins is regarded as the counterpart of the first swarming period of the zoospores of the Saprolegniales. This view derives diplanetism from a more primitive state among the lower fungi, while there is nothing among the algae which corresponds to the phenomenon.

It is further pointed out that some other phenomenon, such as heterogamy, isogamy, and the proliferation of the sporangium, each have a primitive counterpart among the Chytridiales, and can be traced in a natural series to the Saprolegniales and Oomycetes. For some of the phenomena, like the proliferation of the sporangium, there is no counterpart whatever among the algae.—
H. HASSELBRING.

Cultures of Uredineae.—In continuation of the long series of experiments in his cultural work on the Uredineae, the cultures made in 1908 have been reported by ARTHUR.¹⁵ The most interesting result of the work of that year is the discovery that the aecidial stage of Gymnos porangium externum Arthur and Kern (described as new) occurs on a herbaceous perennial, Porteranthus stipulatus (Muhl.) Britton. This is the only case known of the occurrence of an aecidium of Gymnosporangium on a plant outside of the Pomaceae. Another unusual case is that of G. Libocedri (P. Henn.) Kern, whose aecidial stage is shown to be Aecidium Blasdaleanum D. and H., a true aecidium and not belonging to the Roestelia type as in all other Gymnosporangia. Series of cultures with 13 species of rusts gave negative results, and cultures with 23 species confirmed and supplemented work previously reported. The relations of the following species have been worked out for the first time: Puccinia absinthii DC. on Artemisia dracunculoides Pursh has no aecidia, but has pycnidia on the same host; P. macrospora (Peck) Arthur on Carex comosa Boott sown on Smilax hispida Muhl.; P. patruclis Arth. on Carex pratensis Dreej. sown on Agoseris glauca (Pursh) Green; P. cinerea Arth. on Puccinellia airoides (Nutt.) Wats. and Coult. sown on Oxygraphis cymbalaria (Pursh) Prantl; P. Koeleriae Arth. on Koeleria cristata (L.) Pers. sown on Mahonia aquifolium (Pursh) Nutt.; P. alternans Arth. on Bromus Porteri (Coult.) Nash sown on Thalictrum dioicum L.; P. obliterata Arth. on Agropyron biflorum R. and S. sown on Aquilegia canadensis L.; P. Muhlenbergiae Arth, and Holw, on Muhlenbergia glomerata Trin, sown on Callirhoe involucrata (T. and G.) A. Gray; Gymnosporangium Libocedri (P. Henn.) Kern on Libocedrus decurrens Torr. sown on Crataegus Pringlei Sarg.; and G. externum Arth. and Kern on Juniperus virginiana L. sown on Porteranthus stipulatus (Muhl.) Britt.—H. HASSELBRING.

Biology of Gymnosporangium.—Another interesting example showing the slight degree of differentiation existing among some species of parasitic fungi

¹⁵ Arthur, J. C., Cultures of Uredineae in 1908. Mycologia 1:225-256. 1909.